

TECHNICAL DATA SHEET

Sabouraud Dextrose Broth w/Chloramphenicol

Principle

Sabouraud dextrose broth is used for cultivation of yeasts, molds and aciduric bacteria. It is recommended by USP as a medium for microbial limit testing of pharmaceutical products and raw material used in pharmaceutical industries. The medium is prepared in accordance with the harmonized principles of USP. Medium is consisting of meat and casein peptone (1:1) and dextrose monohydrate. The meat and casein peptone provide carbonaceous, nitrogenous compounds, long chain amino acids, vitamins and other essential growth nutrients in addition to that the dextrose monohydrate serves as energy source. The high concentration of dextrose and low pH of medium favor the growth of yeasts and molds and inhibit other contaminating bacteria from pharmaceutical and clinical specimens. In addition to pH the Chloramphenicol provides excess selective for the growth of fungi and yeast.

Use: Used to diagnose fungal infections such as candidiasis, dermatophytosis, and aspergillosis.

Contents*

Ingredients	Gram/Litre
Meat and Casein Peptone	10.000
Dextrose Monohydrate	20.000
Chloramphenicol	0.050
pH at 25°C	5.6 ±0.2

*** Formula adjusted for optimum performance and parameters**

Directions: Dissolve 30.0 grams in 1000 ml distilled water. Sterilize by autoclaving at 15 lbs. pressure (121°C) for 15 min, cool it to 42-45 °C and inoculate test sample aseptically.

Specimens types analyzed

Pharmaceutical samples, clinical samples, etc.

OXFORD LAB FINE CHEM LLP

ISO 9001-2008 Certified Company

Regd Office: Unit no 12, 1st Floor,
Neminath Industrial Estate No.6,
Navghar, Vasai (East), Palghar - 410210.
Maharashtra, INDIA.

Tel: +91 250 2390032 / 2390989 / 2390990
Email: sales@oxfordlabchem.com /
info@oxfordlabchem.com
Web: www.oxfordlabchem.com



Precautions to be taken

These microbial media are intended for the in-vitro use only. All the handling, experiments, storage, and discarding should be performed with the help of skilled and knowledgeable technicians and as per the established guidelines. The material should be disposed only after proper sterilization by autoclaving. Please go through the MSDS of the media to avoid any accidents or in emergency.

Performance and Evaluation

The expected performance of the medium is liable to use as per the direction on the label when stored at optimum conditions and within expiry date.

Quality Control

Appearance	Light beige colored free flowing, homogeneous powder
Reaction of 3.0% solution	5.6 $\pm$ 0.2 at 25 °C
pH	5.40- 5.80 at 25 °C
Color and clarity of ready medium	Light to medium amber, very slightly opalescent without significant precipitate
Growth Promotion properties	Best at $\leq$ 100 CFU at 22-25 °C for 18-72 h
Indicative properties	Optimum at $\leq$ 100 CFU at 22-25 °C for 18-48 h
Negative control	Performed using sterile distilled water

Different Microbial Response

Organism	ATCC	Inoculum CFU	Growth	Incubation temp.	Incubation period
<i>Candida albicans</i>	10231	50-100	Luxurious	22-25 °C	3-5 days
<i>Aspergillus brasiliensis</i>	16404	50-100	Luxurious	22-25 °C	3-5 days
<i>Saccharomyces cerevisiae</i>	9736	50-100	Luxurious	22-25 °C	3-5 days

Storage and Shelf Life: The product is highly hygroscopic; keep the container tightly closed at all times and store it properly as per the conditions mentioned on the label. The declared expiry is valid only when stored as per the conditions mentioned on the label.

Note: Sterilize media immediately after reconstitution.

This document has been produced electronically and it is valid without signature.

www.oxfordlabchem.com

OXFORD LAB FINE CHEM LLP

ISO 9001-2008 Certified Company

Regd Office: Unit no 12, 1st Floor,
Neminath Industrial Estate No.6,
Navghar, Vasai (East), Palghar - 410210.
Maharashtra, INDIA.

Tel: +91 250 2390032 / 2390989 / 2390990
Email: sales@oxfordlabchem.com /
info@oxfordlabchem.com
Web: www.oxfordlabchem.com



Disposal: To avoid the contamination or propagation of any hazardous microbes the used, unusable or modified preparation of this product must be disposed after autoclaving after completion of task.

Reference

1. Atlas, R. M. (2005). *Handbook of media for environmental microbiology*. CRC press.
2. *Difco Manual* (1998). 11th Edition. Difco Laboratories., Division of Becton Dickinson and Company, Sparks, Maryland, USA.
3. Rand, M. C., Arnold E. Greenberg, and Michael J. Taras, (1976), *Standard methods for the examination of water and wastewater*. Prepared and published jointly by American Public Health Association, American Water Works Association, and Water Pollution Control Federation.
4. *The United States Pharmacopoeia*, (2014), The United States Pharmacopoeial Convention. 12601 Twinbrook Parkway, Rockville, MD 20852.

Disclaimer:

The information contained herein in good faith but makes no representations as to its comprehensiveness or accuracy. This document is intended only as a guide to the appropriate precautionary handling of the material by a properly trained person using this product. Individuals receiving the information must exercise their independent judgment in determining its appropriateness for a particular purpose.

Oxford Lab Fine Chem LLP makes no representations or warranties, either express or implied, including without limitation any warranties of merchantability, fitness for a particular purpose with respect to the information set forth herein or the product to which the information refers. Accordingly, Oxford Lab Fine Chem LLP will not be responsible for damages resulting from use of or reliance upon this information.