

TECHNICAL DATA SHEET

Chloramphenicol Yeast Glucose Agar

Principle

Chloramphenicol Yeast Glucose Agar is composed of yeast extract, glucose, chloramphenicol and agar. Yeast extract source of nitrogenous compounds, vitamin B complex and other growth nutrients. Glucose is fermentable sugar. Chloramphenicol inhibits bacterial growth. Agar is solidifying agent. The media is used for enumeration of yeast and molds in milk and milk products.

Use: For isolation & selective enumeration of yeasts and molds in milk & milk products.

Contents*

Ingredients	Gram/Liter
Yeast Extract	5.00
Glucose	20.00
Chloramphenicol	0.10
Agar pH	15.00
at 25°C	6.6 ±0.2

* Formula adjusted for optimum performance and parameters

Directions: Dissolve 40.00 grams in 1000 ml distilled water, boil to dissolve the medium completely. Sterilize by autoclaving at 15 lbs. pressure (121°C) for 15 min, cool it to 42-45 °C and distribute aseptically in sterile petri plates. Ensure complete solidification and inoculate test sample aseptically.

Specimens types analyzed

Milk and milk products and dairy samples etc.

OXFORD LAB FINE CHEM LLP

ISO 9001-2008 Certified Company

Regd Office: Unit no 12, 1st Floor,
Neminath Industrial Estate No.6,
Navghar, Vasai (East), Palghar - 410210.
Maharashtra, INDIA.

Tel: +91 250 2390032 / 2390989 / 2390990
Email: sales@oxfordlabchem.com /
info@oxfordlabchem.com
Web: www.oxfordlabchem.com



Precautions to be taken

All the handling, experiments, storage, and discarding should be performed with the help of skilled and knowledgeable technicians and as per the established guidelines. The material should be disposed only after proper sterilization by autoclaving. Please go through the MSDS of the media to avoid any accidents or in emergency.

Performance and Evaluation

The expected performance of the medium is liable to use as per the direction on the label when stored at optimum conditions and within expiry date.

Quality Control

Appearance	Beige colored free flowing, homogeneous powder
Reaction of 4.00 % solution	6.6 ±0.2 at 25 °C
pH	6.40- 6.80
Gelling	Firm comparable with 1.5% agar gel
Color and clarity of ready medium	Light amber colored slightly opalescent gel
Growth Promotion properties	Best at ≤ 100 CFU at 25-30°C for 2-7 days
Indicative properties	Optimum at ≤ 100 CFU at 32-37 °C for 18-48 h
Negative control	Performed using sterile distilled water

Different Microbial Response: For fungi cultural characteristics observed after an incubation at 25±2°C for 2-5 days.

Organism	ATCC	Inoculum (CFU)	Growth	Recovery
<i>Aspergillus brasiliensis</i>	16404	50-100	Good	≥ 60%
<i>Candida albicans</i>	10231	50-100	Luxuriant	≥ 70%
<i>Escherichia coli</i>	8739	50-100	Inhibited	---
<i>Staphylococcus aureus</i>	25923	50-100	Inhibited	--

This document has been produced electronically and it is valid without signature.

www.oxfordlabchem.com

OXFORD LAB FINE CHEM LLP

ISO 9001-2008 Certified Company

Regd Office: Unit no 12, 1st Floor,
Neminath Industrial Estate No.6,
Navghar, Vasai (East), Palghar - 410210.
Maharashtra, INDIA.

Tel: +91 250 2390032 / 2390989 / 2390990
Email: sales@oxfordlabchem.com /
info@oxfordlabchem.com
Web: www.oxfordlabchem.com



Storage and Shelf Life: The product is highly hygroscopic; keep the container tightly closed at all times and store it properly as per the conditions mentioned on the label. The declared expiry is valid only when stored as per the conditions mentioned on the label. Note: Sterilize media immediately after reconstitution.

Disposal: To avoid the contamination or propagation of any hazardous microbes the used, unusable or modified preparation of this product must be disposed after autoclaving after completion of task.

Reference

1. Atlas, R. M. (2005). Handbook of media for environmental microbiology. CRC press.
2. Marshall, (Ed), (1993), Standard Methods for Examination of Dairy Products, 16th Ed., American Public Health Association, Washington, D. C.

Disclaimer:

The information contained herein in good faith but makes no representations as to its comprehensiveness or accuracy. This document is intended only as a guide to the appropriate precautionary handling of the material by a properly trained person using this product. Individuals receiving the information must exercise their independent judgment in determining its appropriateness for a particular purpose.

Oxford Lab Fine Chem LLP makes no representations or warranties, either express or implied, including without limitation any warranties of merchantability, fitness for a particular purpose with respect to the information set forth herein or the product to which the information refers. Accordingly, Oxford Lab Fine Chem LLP will not be responsible for damages resulting from use of or reliance upon this information.