

TECHNICAL DATA SHEET

Coagulase Mannitol Broth Base

Principle

Coagulase Mannitol Broth Base medium is composed of special peptone, peptone, D-mannitol, sodium chloride and phenol red. Special peptone and peptone, which provide necessary nutrients to the growth of test organisms. D-mannitol is carbon source. Sodium chloride maintains osmotic equilibrium of the medium. The phenol red is pH indicator dye. To detect the coagulase activity the media is fortified with rabbit serum. Coagulase production is dependent on the presence of a fermentable sugar (mannitol) and presence of a protein factor in the brain heart infusion and blood plasma. When mannitol is fermented, the pH of the medium drops. This drop in pH is indicated by the change in color of the phenol red, which turns yellow, and exhibits yellow medium. An opaque broth due to coagulated plasma forms due to growth of coagulase positive organisms. *Staphylococcus epidermidis* is a coagulase negative and mannitol nonfermenting species, does not change the color of the medium. Coagulase negative species may ferment mannitol and produce a yellow color but opacity will not be formed. Production of gas can be determined by placing a small inverted Durham's tube in the medium tube. For the test, use this medium in 2 to 5 ml amounts after adding about 12-15% plasma. Inoculate by adding about 2 drops of test organism and incubate at 37°C and examine after 2-3 hours and also after 4-5 hours of incubation. Un-inoculated control tubes should also be run in parallel with the fermentation tests.

Use: For detection of coagulase production and mannitol fermentation in differentiation of *Staphylococci*.

OXFORD LAB FINE CHEM LLP

ISO 9001-2008 Certified Company

Regd Office: Unit no 12, 1st Floor,
Neminath Industrial Estate No.6,
Navghar, Vasai (East), Palghar - 410210.
Maharashtra, INDIA.

Tel: +91 250 2390032 / 2390989 / 2390990
Email: sales@oxfordlabchem.com /
info@oxfordlabchem.com
Web: www.oxfordlabchem.com



Contents*

	Gram/Litre
Ingredients	
Special Peptone#	375.00
Peptone	10.00
D-Mannitol	10.00
Sodium chloride	5.00
Phenol red	0.025
pH at 25°C	7.4 ±0.2

* Formula adjusted for optimum performance and parameters

Equivalent of Heart Muscle infusion form

Directions: Dissolve 35.00 grams in 1000 ml distilled water. Boil to dissolve the medium completely and distribute in desired. Sterilize by autoclaving at 15 lbs pressure (121 °C) for 15 min, cool it to 42-45 °C and aseptically add sterilized 7-15% rabbit plasm, mix well and distribute in desired and inoculate test sample aseptically.

Specimens types analyzed

Clinical samples, food and dairy samples etc

Precautions to be taken

These microbial media are intended for the in-vitro use only. All the handling, experiments, storage, and discarding should be performed with the help of skilled and knowledgeable technicians and as per the established guidelines. The material should be disposed only after proper sterilization by autoclaving. Please go through the MSDS of the media to avoid any accidents or in emergency.

Performance and Evaluation

The expected performance of the medium is liable to use as per the direction on the label when stored at optimum conditions and within expiry date.

OXFORD LAB FINE CHEM LLP

ISO 9001-2008 Certified Company

Regd Office: Unit no 12, 1st Floor,
Neminath Industrial Estate No.6,
Navghar, Vasai (East), Palghar - 410210.
Maharashtra, INDIA.

Tel: +91 250 2390032 / 2390989 / 2390990
Email: sales@oxfordlabchem.com /
info@oxfordlabchem.com
Web: www.oxfordlabchem.com



Quality Control

Appearance	Light Beige colored free flowing, homogeneous powder
Reaction of 3.50% solution	7.4 ±0.2 at 25 °C
pH	7.20- 7.60
Color and clarity of ready medium	red colored opalescent solution
Growth Promotion properties	Best at ≤ 100 CFU at 32-37 °C for 18-72 h
Indicative properties	Optimum at ≤ 100 CFU at 32-37 °C for 18-48 h
Negative control	Performed using sterile distilled water

Different Microbial Response

Prepare media as per labelled direction. Cultural characteristics observed with added 7-15% v/v sterile pretested coagulase plasma after an incubation at 35-37°C for 18-48 hours.

Organism	ATCC	Growth	Acid from mannitol	Coagulase activity
<i>Staphylococcus aureus</i>	29523	Luxurious	Positive reaction, yellow color	Positive reaction, clot formation
<i>Staphylococcus epidermidis</i>	12228	Inhibited	Negative reaction no color change	Negative reaction, no clot formation
<i>Escherichia coli</i>	8739	Inhibited	Negative reaction no color change	Negative reaction, no clot formation

Storage and Shelf Life: The product is highly hygroscopic; keep the container tightly closed at all times and store it properly as per the conditions mentioned on the label. The declared expiry is valid only when stored as per the conditions mentioned on the label. Note: Sterilize media immediately after reconstitution.

Disposal: To avoid the contamination or propagation of any hazardous microbes the used, unusable or modified preparation of this product must be disposed after autoclaving after completion of task.

This document has been produced electronically and it is valid without signature.

www.oxfordlabchem.com

OXFORD LAB FINE CHEM LLP

ISO 9001-2008 Certified Company

Regd Office: Unit no 12, 1st Floor,
Neminath Industrial Estate No.6,
Navghar, Vasai (East), Palghar - 410210.
Maharashtra, INDIA.

Tel: +91 250 2390032 / 2390989 / 2390990
Email: sales@oxfordlabchem.com /
info@oxfordlabchem.com
Web: www.oxfordlabchem.com



Reference

1. Isenberg, H.D. (2004) Clinical Microbiology Procedure Handbook. Vol. 2, American Society for Microbiology, ASM Press.
2. Jorgensen, J.H., Pfaller, M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock, D.W. (2015) Manual of Clinical Microbiology, 11th Edition. Vol. 1.

Disclaimer:

The information contained herein in good faith but makes no representations as to its comprehensiveness or accuracy. This document is intended only as a guide to the appropriate precautionary handling of the material by a properly trained person using this product. Individuals receiving the information must exercise their independent judgment in determining its appropriateness for a particular purpose.

Oxford Lab Fine Chem LLP makes no representations or warranties, either express or implied, including without limitation any warranties of merchantability, fitness for a particular purpose with respect to the information set forth herein or the product to which the information refers. Accordingly, Oxford Lab Fine Chem LLP will not be responsible for damages resulting from use of or reliance upon this information.